

Enriched Herds And Finite Quantum Groupoids

Thomas Booker and Brian J. Day

April 13, 2010.

Abstract: We describe a finite quantum groupoid associated to any finite $Vect_k$ -enriched herd.

Let $\mathbb{V} = Vect_k$ (finite dimensional k -vector spaces) and let $\mathcal{A}$ be a finite representable k -linear flock (i.e. a finite $\mathbb{V}$ -herd): thus the objects of $\mathcal{A}$ are finite in number, $\mathcal{A}(a, b)$ finite, $\mathcal{A}(a, b)^* \cong \mathcal{A}(b, a)$ chosen (we refer the reader to [1] for the definition of a k -linear flock).

Then the category

$$[\mathcal{A}^{\text{op}} \otimes \mathcal{A}, Vect_k]$$

of finite bimodules over $\mathcal{A}$ is $*$ -autonomous (as shown in [3]).

If also

$$\mathcal{A}(b, \tau(a, c, d))^* \cong \mathcal{A}(a, \tau(b, d, c))$$

then we obtain a functor (i.e. antipode)

$$\begin{aligned} S : \quad \mathcal{H}^{\text{op}} &\longrightarrow \mathcal{H} \\ (x, y) &\mapsto (y, x) \end{aligned}$$

where the objects of $\mathcal{H}$ are the pairs $(a, b) \in \mathcal{A}^{\text{op}} \otimes \mathcal{A}$ and the hom of $\mathcal{H}$ is

$$\mathcal{H}((a, b), (c, d)) = \mathcal{A}(\tau(b, a, c), d) .$$

Moreover, the convolution $[\mathcal{H}, Vect_k]$ with respect to the promonoidal structure on $\mathcal{H}$ given by

$$\begin{aligned} p((a, b), (c, d), (u, v)) &= \mathcal{A}(\tau(d, c, b), v) \otimes \mathcal{A}(u, a) \\ j(u, v) &= \mathcal{A}(u, v) \end{aligned} \tag{1}$$

is $*$ -autonomous for the above S .

There is a prospective Fourier transformation functor $\hat{K}$ on $[\mathcal{H}, Vect_k]$ given by

$$\hat{K}(f)(a, b) = \int_{(x, y) \in \mathcal{H}} f(x, y) \otimes \mathcal{A}(\tau(y, x, a), b)$$

where

$$[\mathcal{H}, Vect_k] \begin{array}{c} \xrightarrow{\hat{K}} \\ \xleftarrow{\tilde{K}} \end{array} [\mathcal{A}^{\text{op}} \otimes \mathcal{A}, Vect_k]$$

with $\hat{K} \dashv \tilde{K}$. The cocontinuous functor $\hat{K}$ is certainly multiplicative and preserves the induced duality on $[\mathcal{H}, Vect_f]$ defined by $f^*(x, y) = f(y, x)^*$. Further, $\hat{K}$ is conservative (as required in [2]).

In fact the Fourier functor $\hat{K}$ is here nothing other than the process of “restriction along h ” where

$$h : \mathcal{A}^{\text{op}} \otimes \mathcal{A} \rightarrow \mathcal{H}$$

is the canonical Kleisli functor mapping $(a, b) \in \mathcal{A}^{\text{op}} \otimes \mathcal{A}$ to $(a, b) \in \mathcal{H}$, which is surjective on objects (see [1] for the details of this construction). The functor $\hat{K} = [h, 1]$ also preserves both the left and right internal homs of the convolution structure on $[\mathcal{H}, \text{Vect}_k]$, this is a straight forward calculation. So $[\mathcal{H}, \text{Vect}_k]$ is a $*$ -autonomous monoidal biclosed category because $[\mathcal{A}^{\text{op}} \otimes \mathcal{A}, \text{Vect}_k]$ is such (and $\hat{K} = [h, 1]$ is conservative). The $*$ -autonomy of $[\mathcal{A}^{\text{op}} \otimes \mathcal{A}, \text{Vect}_k]$ also implies that the domain $\mathcal{H}$ is itself $*$ -autonomous as a promonoidal category over Vect_k , that is, the promultiplication p on $\mathcal{H}$ mentioned above (1) satisfies the cyclic relations given in [2] and [3] for example.

Thus $\hat{K} = [h, 1]$ is a quantum groupoid in the sense of [3] (see also [5]), and there is undoubtedly also a connection with the article by T. Brzeziński and J. Vercruysse: “Bimodule herds”, J. Alg. 321 (2009), 2670-2704. Indeed, the Kleisli functor (or “algebroid”)

$$h : \mathcal{A}^{\text{op}} \otimes \mathcal{A} \rightarrow \mathcal{H}$$

may be related to the Ehresmann Groupoid, even in the more general case where $\mathcal{A}$ is not necessarily finite as a $\mathbb{V}$ -herd.

References

- [1] T. Booker and R.H. Street, *Torsors, herds and flocks*, arXiv:0912.4551 [MATH.CT] Dec 2009 (to appear).
- [2] B.J. Day, *Monoidal functor categories and graphic fourier transforms*, arXiv:math.QA/0612496v1, Dec 2006.
- [3] B.J. Day and R.H Street, *Quantum categories, *-autonomy, and quantum groupoids*, Fields Institute Communications, Volume 43, 2004, pp. 187 - 225.
- [4] G.M. Kelly, *Basic Concepts Of Enriched Category Theory*, London Mathematical Society Lecture Notes Series 64. Cambridge University Press, 1982.
- [5] P. Szlachányi, *Finite Quantum Groupoids And Inclusion Of Finite Type*, Fields Institute Communications, Volume 30, 2001, pp. 393 - 407.

Mathematics Dept., Faculty of Science, Macquarie University, NSW 2109, Australia.

Replies are welcome through Tom Booker (thomas.booker@students.mq.edu.au).